

THE SANTA RITA FAULT ZONE: EVIDENCE FOR LARGE  
MAGNITUDE EARTHQUAKES WITH VERY LONG RECURRENCE  
INTERVALS, BASIN AND RANGE PROVINCE OF  
SOUTHEASTERN ARIZONA

BY PHILIP A. PEARTHREE AND SUSANNA S. CALVO\*

ABSTRACT

**A discontinuous zone of subdued, west-facing fault scarps offset Quaternary alluvium along 58 km of the western piedmont of the Santa Rita Mountains, south of Tucson, Arizona. Scarps trend NE to N-S and are 1 to 6 km basinward from the deeply embayed mountain front. Scarp heights range from 1 to 7 m, and maximum scarp slope angles range from 3° to 9.5°. A 6-m-deep trench excavated in middle Pleistocene and younger deposits exposed a high-angle fault zone with about 3.5 m of total displacement.**

**Estimated ages of faulted and unfaulted surfaces, extent of degradation of piedmont scarps, and stratigraphic relations exposed in the trench suggest an age of about 100 ka for the most recent surface-rupture event. Units having well-developed relict soils of late Pleistocene age and older are faulted, whereas units having weakly to moderately developed soils, of latest Pleistocene and Holocene age, are not offset. Morphologic fault scarp analyses suggest an age of 60 to 100 ka. Earthquake magnitude estimates derived using minimum and maximum seismic source moment calculations range from 6.4 to 7.3.**

**One or more earlier surface-rupture events are implied by significantly higher scarp heights on mid-Pleistocene than on late Pleistocene surfaces. Scarp heights on middle and early (?) Pleistocene surfaces are similar, restricting faulting to post-middle Pleistocene time (<200 to 300 ka). Stratigraphic relationships observed in the trench also suggest two fault-displacement events. Tectonic landform analysis of the mountain front-piedmont area implies several m.y. of inactivity along the range-bounding fault system prior to these late Quaternary displacements. Extremely long recurrent intervals between displacement events and reactivation of faulting after an interval of tectonic quiescence are typical of late Quaternary fault behavior in southeastern Arizona.**

INTRODUCTION

Detailed geomorphic field studies have defined the timing, frequency, and possible magnitudes of late Quaternary surface ruptures on the western piedmont of the Santa Rita Mountains, 30 km south of Tucson, Arizona. This work has implications for the nature of Quaternary faulting in the Basin and Range province of southeastern Arizona and adjacent parts of New Mexico and Sonora, Mexico, where recent studies have delineated widely scattered piedmont fault scarps (Drewes, 1971; Drewes and Thorman, 1980; Morrison *et al.*, 1981; Menges and Pearthree, 1983; Machette *et al.*, 1986). This analysis of the Santa Rita fault zone is useful because it details individual fault behavior in a region where one major historic earthquake has occurred (the Great Sonoran Earthquake of 1887), but where background seismicity is very low (DuBois *et al.*, 1982), and little has been known about recurrence of fault movements.

Fault scarps of the Santa Rita fault zone trend from N-S to NE along about 60

---

\* Present address: Amoco Production Company, Denver, Colorado 80201.




FIG. 1. Location of fault scarps and study area. Numbers indicate portions of the mountain front for which sinuosity values ( $S$ ) (Table 6) were calculated.

km of the southeastern margin of the fault-bounded, north-trending Tucson basin. The scarps are located 1 to 6 km basinward from the topographic mountain front (Figures 1 and 2), range in height from 1 to 7 m, and have maximum slope angles of  $3^{\circ}$  to  $9.5^{\circ}$ . Scarps occur only on surfaces of late Pleistocene age and older.

Tectonic geomorphic analysis, including examination of surficial soils, alluvial stratigraphy, fault scarps, and the topographic mountain front, was used to assess the history of Quaternary surface-rupture faulting events on the Santa Rita piedmont. A soils chronosequence was delineated, and individual surface ages were estimated by comparison with a chronosequence of soils in the southern Rio Grande

valley, New Mexico (Gile *et al.*, 1981). Estimates of the age of most recent faulting are constrained by the age of the oldest surface not offset by faulting and the youngest surface cut by faulting. Greater vertical offset of adjacent older surfaces provides evidence of previous surface faulting. Morphologic scarp-age estimates were made by comparison of the regression line of maximum scarp slope versus log scarp height from the Santa Rita scarps with dated scarps from central Utah (Bucknam and Anderson, 1979). Scarp age was also estimated using a diffusion model of scarp degradation (Nash, 1980; Mayer, 1984). Analysis of a 6-m deep, 37-m-long bulldozer trench crossing the fault provided detailed information on fault zone geometries and evidence for the age of most recent faulting. Field investigations were supplemented by evaluation of the sinuosity and erosional characteristics of the mountain front, parameters which can indicate the relative rate of uplift of the mountain block (Bull and McFadden, 1977).

A zone enveloping the loci of historical seismicity in Arizona extends southeastward from the southern end of the Intermountain seismic belt in northwestern Arizona and may continue through Arizona's southeastern corner to include the epicenter of the 1887 Sonoran earthquake (Sumner, 1976). This event, estimated at magnitude 7.2 to 7.4 (Natali and Sbar, 1982; Herd and McMasters, 1982), produced a 75-km-long (Herd and McMasters, 1982) 0.5 to 4-m-high fault scarp. Northwestern Arizona is the most active part of the zone, with numerous fault scarps cutting Quaternary alluvium and basalt flows, and evidence for extensive late Quaternary offset along some structures (Huntoon, 1977; Hamblin *et al.*, 1981; Menges and Pearthree, 1983). It is not clear, however, if the proposed seismic belt is indeed continuous through the state, as historical seismicity is diffuse (DuBois *et al.*, 1982) and late Quaternary fault scarps uncommon (Menges and Pearthree, 1983) in central Arizona.

#### CHRONOSEQUENCE OF ALLUVIAL GEOMORPHIC SURFACES

Faulted and unfaulted alluvial units deposited on the Santa Rita piedmont provide a means to analyze the Quaternary history of movement on the Santa Rita fault zone. Alluvial units of different ages were defined based on elevation above modern stream channels, degree of surface dissection, surface color, and especially soil-profile development. Ages of units were estimated on the basis of soil-profile development as compared with a dated sequence of soils from the Desert Project of the southern Rio Grande valley of New Mexico (Gile *et al.*, 1981).

*Soil development—Basis for estimating absolute surface ages.* Theoretically, any soil property that increases with age can be used to date a sequence of soils. In order to absolutely date a soils chronosequence, independent absolute age dates are preferable. Unfortunately, we were unable to constrain the absolute ages of surfaces in this area. An alternative and much less precise method for estimating absolute soil ages is to select those soil properties that vary systematically with age in an area and compare them with a dated chronosequence of soils developed from similar parent materials under similar climatic conditions (including the rate of eolian dust fall). Soil properties found to vary consistently with age include redness, maximum amount of clay and iron, and calcic horizon development (see Gile *et al.*, 1966; Bachman and Machette, 1977). These properties were compared with those of a chronosequence of soils defined on the piedmont slopes near Las Cruces, New Mexico (Gile and Grossman, 1979) to estimate ages of surfaces on the Santa Rita piedmont (Tables 1 and 2).

TABLE 1  
SOIL AGE INDICATORS, NONCALCAREOUS PARENT MATERIAL, SANTA RITA PIEDMONT

Unit and Estimated Age (ka)	Maximum Redness (yr unless noted)	Clay		Thickness of Zone of Clay Accumulation (cm)		Iron		Depth to Carbonate Accumulation (cm)	Maximum Stage Development (Bachman and Machette, 1977)
		Maximum % Clay	Increase in %, A to B	Decrease in %, B to C	Maximum % Fe	Increase in %, A to B	Decrease in %, B to C		
Q3b <4	7.5	7	No B horizon	No B horizon	0.59	No B horizon	35	I (weak) or none	
Q3a 4-8	7.5	4	3	2	0.50	0.11	0.16	24-57	II
Q2d 8-20	5	6	5	5	0.43	0.12	0.18	78-148	I (weak)
Q2c 75-130	5 (strong)	22	15	17	0.62	0.24	0.26	141-197	I-III
Q2b-2 200-300	2.5	31	19	28	0.96	0.34	0.74	64	III
Q2b-1 400-750	2.5	44	33	37	1.30	0.55	1.08	141	II-IV
Q2a 1,000-2,000	10 R (weak)	67	42	—	1.27	—	—	Not known	

TABLE 2  
SOIL AGE INDICATORS, PIEDMONT SLOPES NEAR LAS CRUCES, NEW MEXICO [COMPILED FROM GILE AND GROSSMAN (1979) AND GILE ET AL. (1981)]

Unit and Estimated Age (ka)	Clay			Iron			Carbonate		
	Maximum Redness (yr)	Maximum % Clay	Increase in %, A to B	Decrease in %, B to C	Maximum % Fe	Increase in %, A to B	Decrease in %, B to C	Depth to Carbonate Accumulation (cm)	Maximum Stage Development (Bachman and Machette, 1977)
<b>ORGAN</b>									
Mid-to late Holocene 1.1-4.6	5-10	12-18	0-11	1-13	0.8-1.1	0-0.3	0-0.2	25->100	I-II
<b>ISAAC'S RANCH</b>									
Latest Pleistocene 8-15	5	16-28	3-11	4	0.8-0.9	0-0.1	0.1-0.2	13-51	I-III
<b>JORNADA II</b>									
Late Pleistocene 100	2.5-5	28-32	10-19	6-16	0.8-1.0	0-0.3	0-0.3	24-100	III-IV
<b>JORNADA I</b>									
Late Mid-Pleistocene 250-400	2.5-5	33-47	9-20	15-18	1.0-1.3	0.1-0.2	0.3-0.5	18-50	III-V
<b>DONA ANA</b>									
Early to Mid-Pleistocene	10-5	15-74	7-60	—	0.5-2.1	0.1-1.3	—	25-125	IV-V

Several factors affecting soil formation on the Santa Rita and Rio Grande piedmont slopes are similar. Average annual precipitation at soil-profile locations on the Santa Rita piedmont ranges from about 31 to 34 cm (Green and Martin, 1967). The upper piedmont slopes of the Desert Project receive an estimated 35 cm of annual precipitation (Gile and Grossman, 1979, p. 23). More than half the annual precipitation in both areas falls during the summer season. Past climatic variations may have been of a like nature in southern New Mexico and southeastern Arizona, as there is evidence of more effective moisture for soil formation in pre-Holocene soils of both regions (see below, and also Gile and Grossman, 1979, pp. 28–29). Soils used for comparative purposes are developed on parent material dominated by felsic igneous rocks. Rates of eolian dust fall vary substantially across the American Southwest (Machette, 1985), however, adding uncertainty to the proposed soil correlations. Although the variables affecting soil formation are not identical in the two areas, correlation based primarily on the development of soil properties is the best method available to estimate ages of mapped units on the Santa Rita piedmont.

*Discussion of age estimates.* The three youngest units described on the Santa Rita piedmont (Q3b, Q3a, and Q2d, Table 1) exhibit incipient or no clay accumulation and calcic horizon development, and are not offset by faulting. The shallow depth to the top of CaCO<sub>3</sub> accumulation (25 to 60 cm) in soil profiles on Q3 units relative to older soils in the area implies that these soils have developed since the transition to a relatively dry, hot Holocene climate about 8,000 yr ago (Van Devender and Spaulding, 1979). Slightly more clay and redness developed on Q2d imply no great antiquity. However, the greater depth to CaCO<sub>3</sub> accumulation indicates that most of the soil development on Q2d occurred under conditions of more effective soil moisture, probably during the last full glacial (20,000 to 13,000 yr ago) or the relatively wet early Holocene ending about 8,000 yr ago (Van Devender and Spaulding, 1979). The soil parameters for Q2d indicate that it may correlate with the 8,000 to 15,000-yr-old Isaak's Ranch alluvium near Las Cruces (Table 2), consistent with the above arguments.

The four older units in the Santa Rita chronosequence (Table 1) have clay accumulation indicative of substantially greater age, and all have been offset by faulting. Soil parameters for Q2c correlate fairly well with the Jornada II surface of the Las Cruces area, which is about 100,000 yr old (Machette, 1985). The Q2b surfaces are of mid-Pleistocene age (approximately 150 to 750 ka). During this long interval, many separate alluvial units were undoubtedly deposited. Two distinct units were recognized based on soil-profile development. The units differ significantly from each other, but compare generally with the Jornada I surface near Las Cruces. The younger surface, Q2b-2, is probably 200 to 300 ka, while the increase in maximum per cent clay in the soil associated with Q2b-1 suggests that surface is 400 to 750 ka.

Several lines of evidence suggest that Q2a surfaces date to the early Pleistocene and therefore provide a Quaternary-long record of faulting. Q2a soil properties are comparable to well-preserved parts of the Dona Ana surface near Las Cruces, estimated to be early to mid-Pleistocene in age. Additionally, Q2a shares characteristics with other units in southeastern Arizona inferred to be of early Pleistocene age. All are the highest preserved alluvium in their respective basins and have extremely well-developed soils. The soils are developed on relatively thin (<20 m), very coarse-grained, poorly sorted deposits that disconformably overlie finer-grained basin fill deposits and broad bedrock pediments. The degree of soil development

alone argues for the substantial antiquity of these units, and they are the oldest that record the transition from deposition of older basin fill gravels to coarse, poorly sorted Quaternary terrace gravels. Menges and McFadden (1981) postulated that these coarse units were deposited in response to climatic change at or near the beginning of the Quaternary.

### FAULT SCARPS

Topographic profiles of fault scarps surveyed at 61 localities were analyzed to estimate scarp age and frequency of surface displacement. Most profiles were surveyed using an Abney level, tape, and survey rod. Slope measurements were estimated to be accurate to  $\pm 0.5^\circ$ . Profiles were extended beyond the scarps to what appeared to be the pre-faulting surface above and below the scarps. Modification of scarps by gullying and general surface degradation are particular problems in the measurement of old scarps. Areas affected by obvious gullying were avoided, but in many instances surface slopes above and below scarps are not identical, indicating erosional or depositional complications exist. Scarp heights were determined by the difference in elevation between projections of the steepest scarp-slope angle and the "pre-faulting surface" above and below the scarp (after Bucknam and Anderson, 1979).

*Fault scarp-surface age relations.* Fault scarps are preserved only on surfaces of late Pleistocene age (Q2c) and older, indicating the latest fault displacement occurred prior to deposition of Q2d (8 to 20 ka). At several locations, the Q2c surface is displaced less than adjacent scarps on late mid-Pleistocene and older surfaces, implying more than one fault-displacement event. Average scarp heights were determined for the various surfaces to further examine the relation between scarp height and surface age (Table 3). Complex fault splays were excluded because of uncertainty in obtaining total scarp height. Original surface slopes vary only from  $0.5^\circ$  to  $2.5^\circ$ , and therefore do not cause significant increases in scarp height as scarp crests retreat (see Wallace, 1980).

Substantial increase in mean scarp height between Q2c and Q2b surfaces implies at least two distinct fault displacements, one of which occurred prior to deposition of Q2c. Surfaces as young as late mid-Pleistocene (Q2b-2; 200 to 300 ka) record both movements, whereas late Pleistocene Q2c surfaces (75 to 130 ka) have been offset only once. The lack of increase in scarp height between surfaces of early and mid-Pleistocene age suggests that no fault displacement occurred during early to mid-Pleistocene time, unless earlier scarps were completely eroded prior to the late Quaternary movements.

*Morphologic scarp-age estimates.* Two methods were employed to estimate the

TABLE 3  
VARIATIONS IN SCARP HEIGHT WITH SURFACE AGE\*

Surface	Age	No. of Profiles	Range in Height (m)	Mean Height (m)
Q2c	100 ka	11	0.9-3.2	2.21
Q2b	200-750 ka	20	1.5-6.3	4.17
Q2a	1-2 Ma	19	1.0-4.7	3.58
All Q2b-Q2a	—	39	1.0-6.3	3.88

\* Scarps inferred to be the result of complex spatial rupture patterns not included.

age of the Santa Rita fault scarps from their morphology. The first method uses the quantitative relationship between maximum scarp-slope angle and scarp height developed by Bucknam and Anderson (1979). They found that a series of regression lines based on this relationship vary systematically between alluvial scarps of different ages in western Utah. High-stand shoreline scarps of Lake Bonneville provide an absolute time line, from which fault scarp ages can be roughly estimated (see Figure 3).

The second method is a modification of a dating equation based on the diffusion model of hillslope degradation. The dating equation proposed to Nash (1980) states that

$$t_u = t_d(H_u/H_d)^2, \quad (1)$$

where  $t_u$  and  $t_d$  are the ages of the undated and dated scarps, respectively,  $H_u$  is representative height of the undated scarp, and  $H_d$  is the height of a dated scarp with the same maximum slope as the undated scarp. Mayer (1984) modified this method to incorporate the error inherent in estimating the age of scarps of "known" age and the variations in age estimates based on individual profiles of the scarps one seeks to date. He delineated a 10 ka isochron using a least-squares regression to fit a line to the combined scarp-slope versus log of scarp height data from the 14 to 15 ka Lake Bonneville shoreline (Scott *et al.*, 1983) and the early Holocene Drum Mountains scarps (Crone, 1983). Equation (1) is then employed, where  $H_u$  is the



FIG. 2. Aerial view of the Madera Canyon area from the north. Arrows highlight fault scarps that occur on the relatively smooth, light-colored mid-Pleistocene element of Madera Canyon fan (Q2b-1) and slightly darker, late Pleistocene (Q2a) surface to the north. The early Pleistocene element of Madera Canyon fan (Q2a), characterized by much deeper dissection and darker color, is visible south of Q2b-1. The easternmost arrow points to the location of the trench excavated across the fault zone. (Photo courtesy of Peter L. Kresan.)




FIG. 3. Comparison of regression line of maximum scarp slope angle versus log scarp height from the Santa Rita scarp Bonneville shoreline scarps in western Utah (Bucknam and Anderson, 1979). Dashed lines are 95 per cent confidence intervals.

scarp height of an individual profile of the undated scarp. However, the 95 per cent confidence interval on the 10 ka isochron is used to define maximum and minimum values for  $H_d$ . Using these two values in the age estimation equation, minimum and maximum age estimates are produced for each scarp profile. All age range estimates for the individual profiles of the undated scarp are compiled in a histogram, in which the mode and the scatter in the age estimates can be evaluated (Figure 4).

The dating methods employed depend on Lake Bonneville shoreline scarps for absolute age calibration; therefore, variables that control rates of scarp degradation must be considered before age estimates of the Santa Rita fault scarps can be attempted. The climates of western Utah and southeastern Arizona are semi-arid. More precipitation occurs as intense, runoff-producing summer thunderstorms in southeastern Arizona (Rumney, 1968), but freeze-thaw activity is more prevalent in Utah. Scarps on the Santa Rita piedmont occur on unconsolidated, cobbly alluvium similar to that described by Bucknam and Anderson (1979) in Utah. Some cohesion is provided by pedogenic clay and carbonate accumulations, and this may tend to retard scarp degradation. Changes in precipitation and vegetation between glacial and interglacial intervals have likely resulted in changes in rates of scarp degradation in western Utah and southeastern Arizona. Increased precipitation during glacial intervals would increase the power available to degrade scarps. However, this is probably more than offset by increased vegetative density, resulting in lower rates of scarp degradation than determined for the Holocene (Machette, cited in Pierce, 1981). Scarp aspect is another important variable, particularly when comparing north- and south-facing scarps (K. L. Pierce, 1981). All of the Santa Rita scarps face generally west; the effect of this aspect is uncertain.




FIG. 4. Compilation of diffusion-based age-range estimates for scarps on surfaces of all ages (after Mayer, 1984), probably includes composite scarps. *Shaded area and inset* include age-range estimates of probable single event scarps on the late Pleistocene Q2c surface only.

Bucknam and Anderson (1979) confined their measurements to scarps interpreted to have been produced by one event or several events closely spaced in time relative to the interval since the last event. Most of the fault scarps surveyed on the Santa Rita piedmont are probably composite, but examination of individual scarp profiles did not consistently reveal segmented profiles by which discrete events might be discerned (Wallace, 1977). Therefore, the composite scarps appear to fall into the latter category described by Bucknam and Anderson (1979). When comparison is made between composite scarps and single-rupture scarps, composite scarp heights are exaggerated and the age of the most recent event is overestimated (Mayer, 1984). In summary, the factors controlling degradation of the Santa Rita scarps are similar to those of Utah, but the many possible variations preclude a high level of precision in age estimates.

Regression lines obtained from the Santa Rita scarps and from Lake Bonneville shoreline scarps are shown in Figure 3. Correlations of the Santa Rita regression lines were vastly inferior to those of the scarps in Utah. Using scarp profiles from all age surfaces, the coefficient of determination ( $R^2$ ) is only 0.38. Regressions were also run on subsets of the data, grouped both according to age of surface and geographical location, with little improvement in  $R^2$  values. These poor correlations may be due to scarp antiquity, dissection, and/or the inclusion of composite scarps. They clearly increase the uncertainty of age estimates based on comparison with the regression lines from Utah. However, the Santa Rita scarps appear to be distinctly older than the Bonneville shoreline scarps, i.e., at least late Pleistocene in age.

Diffusion-based scarp-age estimates (Figure 4) suggest an age of 60 to 100 ka for the Santa Rita scarps. The modal value obtained for age estimates of all scarps is slightly less than 100,000 yr old (Figure 4a). Analysis of probable single-event scarps on the late Pleistocene Q2c surface indicates an age of 60 to 75 ka. If, however, the rate of scarp degradation was lower during the late Pleistocene than during the Holocene, then all morphologic scarp-age estimates may be too young.

## TRENCH STRUCTURE AND STRATIGRAPHY

A 6 m × 37 m bulldozer trench excavated across the Santa Rita fault zone about  $\frac{1}{2}$  km northeast of the Madera Canyon fan (see Figure 2) provided a detailed exposure of the shear zone in the Quaternary alluvium. The late Pleistocene Q2c surface is offset 2 to 3 m in this area, although no scarp exists at the surface of the trench site because it was dug in a small arroyo for practical and environmental reasons.

Mapping of the trench walls (Figure 5) revealed a sequence of seven thin Pleistocene alluvial deposits and two buried soils, and a thin, discontinuous late (?) Holocene deposit. All but the youngest two deposits were offset across a zone of steeply dipping normal faults during one or two faulting events. The sediments, ranging from poorly sorted conglomerate to massive silt, are characteristic of fluvial deposition on the piedmont in response to climatic variations that changed the discharge of water and sediment from the hillslopes of the Santa Rita Mountains (see Table 4 for description of units). Two depositional hiatuses in the sequence are indicated by buried paleosols developed on units I and II. A surficial soil has formed across the faulted and subsequently beveled alluvial sequence.

*Structure.* The 3.3 to 3.5 m total offset measured on units I to IV is distributed across a 3.5-m-wide zone of 70° to 90° westward-dipping normal faults. Thinning, passive folding, tilting, general disruption of bedding, and some drag structures are associated with the faulting. In plan view, the faults anastomose along strike. The 2-m-wide step in the trench excavation provided an excellent exposure of their pattern. The main fault plane branches out across the step to include five faults spread across 90 cm of very disrupted sediments.

*Age of the most recent faulting event.* Stratigraphic evidence in the trench suggests the most recent fault rupture occurred ~100 ka, in fair agreement with morphologic scarp-age estimates of 60 to 100 ka. The oldest unfaulted unit exposed in the trench (unit VII) provides a minimum age for the latest fault movement. Characteristics of the soil developed on unit VII (Table 5) correlate fairly well with those of the Q2d piedmont unit (Table 1), estimated to be 10 to 20 ka. At least part of the deposition of unit VII appears to be directly related to faulting, however, and predates the climatically caused deposition of the Q2d unit elsewhere on the piedmont. The lower part of the deposit at the main fault trace contains crude cobble layers and abundant gravel, which probably represent debris deposited at the base of the scarp. Unfortunately, any additional scarp-related features at the top of the trench section were eroded along with the scarp.

The absence of the late Pleistocene Q2c soil in the downthrown block of the exposure, even though the faulted Q2c geomorphic surface occurs adjacent to the trench site, suggests that the latest faulting event occurred shortly after stabilization of the Q2c surface, before substantial soil-profile development had occurred. In the trench, unit VI is tentatively correlated with Q2c because it is the youngest faulted unit; however, unit VI displays very little evidence of soil development. Substantial erosion of unit VI does not appear to have taken place, as units VI to VII contact is quite subtle. It appears, therefore, that faulting occurred shortly after deposition of the Q2c unit, about 100 ka.

*Evidence for an earlier faulting event.* Several lines of evidence suggest that two faulting events are recorded in the trench stratigraphy. Units I through IV are all vertically offset about 3.4 m, and their thicknesses do not increase on the downthrown side of the fault, indicating all faulting postdates deposition of unit IV. The




FIG. 5. Interpretation on trench excavated across the Santa Rita fault zone just north of Madera Canyon fan. Upper trench wall is set back from lower wall along 2.5- to 3-m step. Main shear zone in upper wall splays across the step to include five shear planes in lower wall, as indicated by arrows. See Table 5 for stratigraphic descriptions.

TABLE 4  
DESCRIPTION OF ALLUVIAL UNITS MAPPED IN THE TRENCH  
EXPOSURE\*

Unit	Thickness (cm)	Description
Paleosol 3		
VII	70-100	Silty sand and gravel. Massive, moderately sorted, slightly hard, lower boundary clear to gradual.
VI	70-100	Silty sand and gravel. Massive, moderately sorted, hard, lower boundary gradual to diffuse.
V	80(?) - 180	Cobbly fluvial gravel. Moderately well bedded, with sandy beds and cobbly beds, cut and fill structures. Locally well-developed cross-bedding in sandy layers, moderately well to poorly sorted, fairly loose to slightly hard; near main shear zone the inclined sandy gravel layers 4- to 5-cm-thick have well-developed red (2.5 yr 4/8) 1- to 2-cm-thick B-bands; lower boundary abrupt, wavy, locally eroded into or through unit IV.
IV	0-64	Silt. Massive, well sorted, very hard, vesicular, occasional slightly gravelly crude layers 2- to 3-cm-thick, lower boundary abrupt, wavy, locally eroded into unit III.
Paleosol 2		
III	45	Gravelly silty sand. Moderately well sorted, slightly hard to hard, some cross-bedding, lower boundary gradual to diffuse.
II	85-115	Cobbly gravel and silty sand. Moderately well-sorted, loose to slightly hard, weak imbrication, discontinuous slightly hard silt layers 10- to 20-cm-thick, basal slightly hard silty sand layers 10- to 15-cm-thick, lower boundary abrupt and straight.
Paleosol 1		
I	280+	Very cobbly boulder gravel. Massive, very poorly sorted, moderately well-to-well consolidated, lower boundary not exposed.

\* See Table 5 for Paleosol descriptions.

thickness of unit V measured on the downthrown block increases from 1.2 to 1.8 m downslope away from the main shear zone. Projection of the top of unit V from the eastern end of the trench suggests it was about 0.8-m-thick immediately upslope from the fault zone, although the original thickness of unit V may not be preserved

in the upthrown block. The fact that unit V thickens across the fault zone implies that it was deposited over preexisting topography, probably a fault scarp.

Deposition of unit V shortly after a faulting event would explain the complex and disrupted sedimentary structures in unit V, located within  $1\frac{1}{2}$  m downslope from the main fault shear. These may be debris deposited at the base of a low scarp that was subsequently faulted during the most recent displacement. Additionally, several of the faults west of the main shear zone offset units I to IV but not unit V. These relations suggest that offset during an earlier event involving units I to IV was distributed across a 3.5-m-wide zone of faults, while almost all offset during the most recent event occurred on the main shear zone. Vertical displacement attributable to the most recent faulting event is about 2.2 m, assuming the reconstructed top of unit V was originally uninterrupted across the fault zone.

Comparison of offset of trench units with surface offset calculated from scarp profiles on Q2c and Q2b surfaces near the trench supports the hypothesis that two surface ruptures have occurred. The average surface offset of the Q2c unit (2.3 m) is about the same as that postulated for the most recent event in the trench. Q2b-1 surface offset of about 3.2 m on the Madera Canyon fan, 0.5 km to the south, is very similar to total offset determined in the trench. These relations suggest that both the trench section and the mid-Pleistocene Q2b-1 surface record two faulting events, and that the late Pleistocene Q2c surface records only the most recent event.

#### ESTIMATED MAGNITUDE OF PALEOSEISMIC EVENT(S)

Correlation between seismic source moment ( $M_0$ ) and earthquake magnitude ( $M$ ) allows estimation of earthquake sizes associated with the scarp-producing events.  $M_0$  is the product of average fault dislocation ( $\langle u \rangle$ ), the shear modulus ( $\mu$ ), and fault plane area ( $A$ )

$$M_0 = \langle u \rangle \mu A \quad (2)$$

(Brune, 1968). Vertical surface offset and length of scarps can be measured, but the amount of offset may be uncertain if a scarp actually represents multiple events, and total length of a prehistoric fault may be obscured by subsequent erosion or deposition across various scarp segments. Segments of the fault might rupture separately, although within the limits of the timing data we did not find evidence of this. Also, estimated average surface displacement is only a proxy for average fault dislocation, which is not determinable for prehistoric earthquakes. The value used for the shear modulus is  $3.3 \times 10^{11}$  dyne/cm<sup>2</sup> (Brune, 1968; Hanks and Kanamori, 1979). Total depth of brittle failure is probably 10 to 15 km, as microearthquakes detected during a study of the Pitaycachi fault (Natali and Sbar, 1982) were no deeper than 16 km. Dip angles of 60° to 70° have been observed for range-bounding and bedrock normal faults in the southern Basin and Range province (Menges, 1981, unpublished data; Natali and Sbar, 1982). Fault geometries at depth are not constrained; if faults flatten at depth, fault plane area would be greater than assumed here. The calculated value of  $M_0$  is used to estimate paleoearthquake magnitude through a moment-magnitude scale (Hanks and Kanamori, 1979)

$$M = \frac{2}{3} \log M_0 - 10.7. \quad (3)$$

TABLE 5  
SELECTED SOIL CHARACTERISTICS, SURFICIAL AND BURIED SOILS IN TRENCH SECTION

Parent Material (Alluvial Units)	Maximum Redness	Thickness (cm) Zone of Clay Accumulation	Wet Consistence B2t <sub>h</sub> Horizon	Texture B2t <sub>h</sub> Horizon	Depth to Carbonate Accumulation (cm)	Maximum Stage CaCO <sub>3</sub> Development (Bechman and Machette, 1977)	Correlative Piedmont Surficial Soil
3a VI, VII, surface, down-thrown block	5 yr 4/6	91	Sticky, plastic	Gravelly, silty clay loam	76	I-II	Q2d
3b V-II, surface, up- thrown block	5 yr 4/6	78	Sticky, plastic	Very gravelly, sandy clay loam	85	I-II	Q2c or Q2d
2 III, II, buried, down-block	5 yr 4/8	52	Sticky, slightly plastic	Very gravelly, sandy clay loam	None	—	—
1 I, buried, upthrown block	2.5 yr 4/8	280+	Very sticky, very plastic	Cobbly, clay	43	II	Q2b-1

Combining regional structural characteristics with specific offset and fault length data collected from the Santa Rita fault scarps, a range of possible paleoearthquake magnitudes was calculated for the fault system. A minimum value of  $M = 6.4$  is derived using 1.2 m average offset, 10 km length (total length of the northernmost scarp segment), and a fault plane dipping  $70^\circ$  extending to 10 km depth. The maximum value of  $M = 7.3$  is derived with 3.5 m average offset on a fault plane dipping  $60^\circ$ , extending to 15 km depth, along the entire 58 km length from the scarps. A more realistic value might be  $M = 6.9$ , which could be generated by an average of 2 m offset along the 35 km of the fault north from the Madera Fan area.

#### PLIO-QUATERNARY ACTIVITY ON THE SANTA RITA FAULT ZONE

The Santa Rita Mountain front was analyzed using landscape parameters which can indicate relative rates of vertical tectonic activity at mountain fronts (Bull and McFadden, 1977). Soil, stratigraphic, and scarp morphology studies restrict times of Quaternary faulting on the Santa Rita piedmont to late mid- and late Pleistocene. Analysis of the tectonic landforms of the mountain front, which extends our analysis of the faulting history back probably through the Pliocene, suggests that relative tectonic inactivity along the range front began at least 3 to 5 m.y. ago.

Calculation of landscape parameters allows quantification of the relative rates of base-level processes. Mountain-front sinuosity ( $S$ ) and valley floor width-valley height ratios ( $Vf$ ) compare the rate of uplift along the range-bounding structures and rates of fluvial erosion. Sinuosity ( $S$ ) is defined as the ratio of the length along the actual bedrock/alluvium contact to the overall length of the mountain front, both measured from topographic maps. High  $S$  values indicate the mountain front has been erosionally embayed from its originally more linear, fault-bounded configuration.  $Vf$  ratios for drainages along the mountain front can be used as an index of whether streams are actively downcutting or primarily eroding laterally.  $Vf$  is the ratio of valley floor width to the average elevation of the left and right divides above the stream channel. Large  $Vf$  ratios, indicating lateral stream planation, is dominant relative to downcutting, a situation that occurs when uplift rates are low or nonexistent along range-front structures.

The mountain front was divided into five lithologically distinct parts (Figure 1) for which sinuosity and representative  $Vf$  ratios were calculated from 1:24,000 scale topographic maps.  $Vf$  ratios were determined at one-tenth (0.1) the distance of basin length up from the mountain front for moderate-sized drainage basins. The valley floor was defined as the gently sloping surface between steep hillslope walls.

Resulting values of  $S$  and  $Vf$  (Table 6) are characteristic of tectonically inactive fronts (Bull and McFadden, 1977). Variations in the values of  $S$  and  $Vf$  for the Santa Rita Mountains are related to differing resistance to erosion of the lithologic types and relative sizes drainage basins present along each mountain-front segment. Tectonic inactivity along the western Santa Rita front is also suggested by the presence of inselbergs and a broad pediment mantled by thin alluvial fan deposits. A regional depth-to-bedrock survey (Oppenheimer and Sumner, 1980) indicates that shallowly buried ( $\approx 120$  m) bedrock benches occur between the outer inselbergs and mountain front.

The length of time elapsed since substantial faulting has occurred along the Santa Rita Mountains can be estimated by the time required to develop the observed bedrock pediment. Mayer (1979) estimated that the Mogollon Rim in east central

TABLE 6  
 LANDSCAPE PARAMETERS CALCULATED FOR THE WESTERN  
 SANTA RITA MOUNTAIN FRONT (AFTER BULL AND  
 MCFADDEN, 1977)

Mountain-Front Segment	Lithology	S	Basin Length* (km)	V <sub>f</sub>
1	K quartz monzonite	3.9	5.5	57.5
			1.6	58.3
2	Paleozoic and K sediments	3.5	3.2	22.7
			6.4	72.4
3	p <sub>e</sub> granodiorite	3.6	4.2	19.2
			2.4	40.0
			3.1	30.0
4	Tertiary volcanics; K sediments; p <sub>e</sub> schist; complex faults	3.1	5.0	3.5
			5.5	26.3
5	mid-K granodiorite; quartz monzonite; quartz veins	2.4	5.5	5.3

\* Long drainage basins ( $\geq 5.0$  km) also tend to be very wide. Thus, their area is proportionately much larger than shorter basins.

Arizona has retreated at a rate of 0.35 to 0.9 km/m.y. during the late Cenozoic. Denudation rates of  $\sim 1$  km/m.y. have been suggested for coarse-grained plutonic rocks of north-central Nevada and the White Mountains of California (Marchand, 1971; Wallace, 1978). Pediment width in the study area can only be estimated as it is mantled with alluvium. A minimum estimate of 2 to 3.8 km is obtained from the distance between the outermost bedrock exposures and the mountain-front escarpment. The measured distance of the fault scarps from the mountain front (2 to 6 km; mean of  $\sim 4$  km) was used as a maximum pediment width. Using maximum and minimum pedimentation rates, these data imply a 2 to 10 m.y. interval since uplift rates exceeded rates of mountain-front retreat.

#### REGIONAL NEOTECTONIC IMPLICATIONS

Regional evidence is consistent with the late Cenozoic period of quiescence along the Santa Rita front. Menges and McFadden (1981) concluded that major faulting of the Basin-Range disturbance in southeastern Arizona occurred between 13 to 10 Ma (Scarborough and Peirce, 1978; Shafiqullah *et al.*, 1978) and 3 to 6 Ma. Their conclusion was based on paleomagnetic stratigraphy in upper basin-fill deposits of Sonoita Creek basin (east of the Santa Rita Mountains) and the presence of dissected bedrock pediments 2- to 7-km wide, overlain by thin Quaternary deposits, in several basins of southeastern Arizona.

Field studies have been conducted on other late Quaternary fault scarps in southeastern Arizona and adjacent New Mexico and Sonora, Mexico (Menges and Pearthree, 1983; Machette *et al.*, 1986). These studies have demonstrated that late Quaternary faulting in this region has been concentrated in a belt near the Arizona-New Mexico border. Several fault scarps, including the Santa Rita scarps, occur peripheral to this belt. Recurrence intervals between movements on individual faults, where they can be inferred, are consistently on the order of  $10^5$  yr. Pediments

of varying widths exist along all mountain fronts associated with late Quaternary fault scarps, implying that reactivation after a period of tectonic quiescence and/or very low long-term rates of faulting are characteristic of the region.

The long period of tectonic inactivity in southeastern Arizona since the main phase of the Basin-Range disturbance (5 to 10 m.y.) raises questions about the implications of late Quaternary faulting along inferred Basin-Range structures. This long period of relative tectonic quiescence, with a few faults being active occasionally, may be a final phase of the Basin-Range disturbance (e.g., Shafiqullah *et al.*, 1978). Alternatively, Quaternary activity on Basin-Range faults can be interpreted as evidence for transition to a new, weakly extensional tectonic regime that has developed after cessation of the Basin-Range disturbance. A low regional strain rate is implied by the long faulting recurrence intervals ( $\sim 10^5$  yr) inferred for all faults in southeastern Arizona.

### SUMMARY

Late Quaternary activity along the Santa Rita fault zone has been characterized by infrequent but large-magnitude, surface-rupture faulting events. Studies of soils, stratigraphy, and scarp morphology indicate that Quaternary faulting has been restricted to the last 200 to 300 ka. One surface-rupture event is inferred to have occurred between about 130 to 300 ka and another occurred about 60 to 100 ka. Magnitude estimates for these events range from 6.4 to 7.3, with 6.9 considered most reasonable. Geomorphic analysis of the Santa Rita mountain front indicates that it was tectonically quiescent for at least several million years prior to deposition of early and mid-Pleistocene alluvial units.

Recurrence intervals between surface-faulting events on the order of  $10^5$  yr and reinitiation of faulting after an interval of tectonic quiescence have been inferred for other Quaternary faults studied in southeastern Arizona and adjacent New Mexico and Sonora, Mexico. Long recurrence intervals and the limited extent of Quaternary faulting in the region imply a relatively weak regional extensional stress that has reactivated some favorably oriented older structures. This reactivation is evidence either of a late pulse of activity related to the Basin-Range event, or a new, weakly extensional tectonic regime that has developed during the Quaternary.

### ACKNOWLEDGMENTS

We would like to thank our advisor, Bill Bull, for imparting enough of his knowledge of tectonic geomorphology to permit us to undertake and complete this study. We also thank Larry Mayer, Dave Hendricks, Pete Knuepfer, Les McFadden, Chris Menges, and Marc Sbar for technical advice and many helpful discussions. Review by Marie Slezak Peartree, Carl Glass, and M. N. Machette improved this manuscript substantially. Funding was provided by National Science Foundation Grants EAR 78-03648 and EAR 80-08212 ("Seismic and Geomorphic Studies of Quaternary Tectonism in the Arizona Seismic Belt").

### REFERENCES

- Bachman, G. O. and M. N. Machette (1977). Calcic soils and calcretes in the southwestern United States, *U.S. Geol. Surv., Open-File Rept. 77-794*, 163 pp.
- Brune, J. N. (1968). Seismic moment, seismicity and rate of slip along major fault zones, *J. Geophys. Res.* **75**, 777-784.
- Bucknam, R. C. and R. E. Anderson (1979). Estimation of fault scarp ages from a scarp-height-slope-angle relationship, *Geology* **7**, 11-14.

- Bull, W. B. and L. D. McFadden (1977). Tectonic geomorphology north and south of the Garlock fault, California, in *Geomorphology in Arid Regions*, Doehring, D. O., Editor, *Proceedings, Eighth Annual Geomorphology Symposium*, SUNY, Binghamton, New York, 115–138.
- Crone, A. J. (1983). Field trip 5: amount of displacement and estimated age of a Holocene surface faulting event, eastern Great Basin, Milliard County, Utah, in *Geologic Excursions in Neotectonics and Engineering Geology in Utah; Guidebook, Part IV*, Gurgel, K. D., Editor, *Special Studies—Utah Geological and Mineral Survey* **62**, 49–55.
- Drewes, H. (1971). Geologic map of the Sahuarita Quadrangle, southeast of Tucson, Pima Co., Arizona, U.S. Geol. Surv. Misc. Invest. Series Map I-613, scale 1:48,000.
- Drewes, H. and C. H. Thorman (1980). Geologic map of the Cotton City quadrangle and the adjacent part of the Vanar quadrangle, Hidalgo County, New Mexico, U.S. Geol. Surv. Misc. Invest. Series Map I-1221, scale 1:24,000.
- DuBois, S. M., A. W. Smith, N. K. Nye, and T. A. Nowak (1982). Arizona Earthquakes, 1776–1980, *Arizona Bureau of Geology and Mineral Technology Bulletin* **193**, 456 pp.
- Gile, L. H. and R. B. Grossman (1979). The Desert Project Soil Monograph, U.S. Department of Agriculture, Soil Conservation Service, 984 pp.
- Gile, L. H., F. F. Peterson, and R. B. Grossman (1966). Morphological and genetic sequences of carbonate accumulation in desert soils, *Soil Sci.* **101**, 347–360.
- Gile, L. H., J. W. Hawley, and R. B. Grossman (1981). Soils and geomorphology in the Basin and Range area of southern New Mexico—Guidebook to the Desert Project, *New Mexico Bur. Mines Min. Res., Memoir* **39**, 218 pp.
- Green, C. R. and S. C. Martin (1967). An evaluation of precipitation, vegetation and related factors on the Santa Rita Experimental Range, University of Arizona, Institute of Atmosphere and Physics, *Technical Reports on the Meteorology and Climatology of Arid Regions* **17**, 82 pp.
- Hamblin, W. K., P. E. Damon, and W. B. Bull (1981). Estimates of vertical crustal strain rates along the western margin of the Colorado Plateau, *Geology* **9**, 293–298.
- Hanks, T. C. and H. Kanamori (1979). A moment magnitude scale, *J. Geophys. Res.* **84**, 2348–2350.
- Herd, D. G. and C. R. McMasters (1982). Surface faulting in the Sonora, Mexico earthquake of 1887, *Geol. Soc. Am. Abstracts with Programs* **14**, 172.
- Huntoon, P. W. (1977). Holocene faulting in the western Grand Canyon, Arizona, *Geol. Soc. Am. Bull.* **88**, 1619–1622.
- Machette, M. N. (1985). Calcic soils of the American Southwest, *Geol. Soc. Am. Special Paper* **203**, 1–21.
- Machette, M. N., S. F. Personius, C. M. Menges, and P. A. Pearthree (1986). Quaternary and Pliocene faults in the Silver City and Douglas 1° × 2° quadrangles, U.S. Geol. Surv. Map MF 1465-C, scale 1:250,000.
- Marchand, D. E. (1971). Rates and modes of denudation, White Mountains, eastern California, *Am. J. Sci.* **262**, 417–435.
- Mayer, L. (1979). Evolution of the Mogollon rim in central Arizona, *Tectonophysics* **61**, 49–62.
- Mayer, L. (1984). Dating Quaternary fault scarps formed in alluvium using morphologic parameters, *Quat. Res.* **22**, 300–313.
- Menges, C. M. and L. D. McFadden (1981). Evidence for a latest-Miocene to Pliocene transition from Basin-Range tectonic to post-tectonic landscape evolution in southeastern Arizona, *Arizona Geol. Soc. Dig.* **13**, 151–160.
- Menges, C. M. and P. A. Pearthree (1983). Map of neotectonic (latest Pliocene/Quaternary) deformation in Arizona, *Arizona Bur. Geol. Min. Tech. Open-File Rept.* **83-22**, scale 1:500,000.
- Morrison, R. B., C. M. Menges, and L. K. Lepley (1981). Neotectonic maps of Arizona, *Arizona Geol. Soc. Dig.* **13**, 179–183.
- Nash, D. B. (1980). Morphologic dating of degraded normal fault scarps, *J. Geol.* **88**, 353–360.
- Natali, S. G. and M. L. Sbar (1982). Seismicity in the epicentral region of the 1887 northeastern Sonora earthquake, Mexico, *Bull. Seism. Soc. Am.* **72**, 181–196.
- Oppenheimer, J. M. and J. S. Sumner (1980). Depth-to-bedrock map, Basin and Range province, Arizona, Laboratory of Geophysics, University of Arizona, Tucson, Arizona, scale 1:1,000,000.
- Pierce, K. L. (1981). Quaternary dating and neotectonics, *U.S. Geol. Surv., Open-File Rept.* **81-833**, 154–155.
- Rumney, G. R. (1968). *Climatology and the World's Climates*, MacMillan Co., New York, 403–404.
- Scarborough, R. B. and H. W. Peirce (1978). Late Cenozoic basins of Arizona, *New Mexico Geol. Soc. Guidebook* **29**, 253–259.
- Scott, W. E., W. D. McCoy, R. R. Shroba, and R. Meyer (1983). Reinterpretation of the exposed record of the last two cycles of Lake Bonneville, western United States, *Quat. Res.* **20**, 261–285.

- Shafiqullah, M., P. E. Damon, D. J. Lynch, P. H. Kuck, and W. A. Rehrig (1978). Mid-Tertiary magmatism in southeastern Arizona, *New Mexico Geol. Soc. Guidebook* **29**, 231-241.
- Sumner, J. S. (1976). Earthquakes in Arizona, *Field Notes from the Arizona Bureau of Mines* **6**, no. 1, 1-2.
- Van Devender, T. R. and W. G. Spaulding (1979). Development of vegetation and climate in the southwestern United States, *Science* **204**, 701-710.
- Wallace, R. E. (1977). Profiles and ages of young fault scarps, north-central Nevada, *Geol. Soc. Am. Bull.* **88**, 1267-1281.
- Wallace, R. E. (1978). Geometry and rates of change of fault-generated range fronts, north-central Nevada, *U.S. Geol. Surv. J. Res.* **6-6**, 637-650.
- Wallace, R. E. (1980). Nomograms for estimating components of fault displacement from measured height of fault scarp, *Assoc. Eng. Geol. Bull.* **17**, 39-45.

DEPARTMENT OF GEOSCIENCES  
UNIVERSITY OF ARIZONA  
TUCSON, ARIZONA 85721

Manuscript received 24 May 1985